


Complimentary Resources to Support Your Care Team and Patients

Streamline your team and patient experience

Chart Sticker

Description:
A sticker affixed to the patient chart


For use by:



- ① **Nursing staff:** Local anesthetic utilization window reminder for PACU, and floor nurses
- ② **Billers:** Procedure date and dose tracking

ASC C9290 OR Baggie

Stores vials, wristband, and ASC OR log sticker



- ① **Nursing staff:** Conveniently store OR items together for the procedure
- ② **Billers:** ASC OR log stickers to support billing documentation requirements

Syringe Label


A sticker placed on syringes of EXPAREL ready for infiltration



- ① **Nursing staff:** Time-stamped labels to distinguish syringes of EXPAREL being used in a procedure

ASC OR Log Stickers


A sticker for the OR log to document the amount of EXPAREL that is used AND unused in a procedure



- ① **Nursing staff:** Used and unused dose to be provided by the surgeon
- ② **Billers:** Both used and unused dose must be noted. Billing is per mg, not per mL
- ③ **Billers:** HCPCS code C9290 is used to bill for EXPAREL use in ASCs

Patient Wristband


Wristband with EXPAREL administration timing



- ① **Nursing staff:** Administration reminder
- ② **Nursing staff:** Verification indicating when a patient may receive additional local anesthetic
- ③ **Patients:** Awareness they have received an opioid-minimizing protocol

Patient Discharge Instructions

Patient education resource outlining how to manage postsurgical pain after discharge



- ① **Nursing staff:** Details the postsurgical care plan
- ② **Patients:** Personalized take-home treatment and recovery information

*Also available in Spanish

ASC=ambulatory surgery center; HCPCS=Healthcare Common Procedure Coding System; OMFS=oral and maxillofacial surgery; OR=operating room; PACU=postanesthesia care unit.



ORDER COMPLIMENTARY ITEMS

Contact your Pacira representative or scan the QR code with your mobile device.

Please see Important Safety Information on next page.
Full Prescribing Information is also available at www.EXPAREL.com.



Indication

EXPAREL is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia.

Adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via infiltration were nausea, constipation, and vomiting; adverse reactions reported in adults with an incidence greater than or equal to 10% following EXPAREL administration via interscalene brachial plexus nerve block were nausea, pyrexia, and constipation.

Adverse reactions with an incidence greater than or equal to 10% following EXPAREL administration via infiltration in pediatric patients six to less than 17 years of age were nausea, vomiting, constipation, hypotension, anemia, muscle twitching, vision blurred, pruritus, and tachycardia.

If EXPAREL and other non-bupivacaine local anesthetics, including lidocaine, are administered at the same site, there may be an immediate release of bupivacaine from EXPAREL. Therefore, EXPAREL may be administered to the same site 20 minutes after injecting lidocaine.

EXPAREL is not recommended to be used in the following patient populations: patients <6 years old for infiltration, patients younger than 18 years old for interscalene brachial plexus nerve block, and/or pregnant patients.

Because amide-type local anesthetics, such as bupivacaine, are metabolized by the liver, EXPAREL should be used cautiously in patients with hepatic disease.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL.

EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use.

The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression.

Cardiovascular System Reactions: Toxic blood concentrations depress cardiac conductivity and excitability, which may lead to dysrhythmias, sometimes leading to death.

Allergic Reactions: Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients.

Chondrolysis: There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

Methemoglobinemia: Cases of methemoglobinemia have been reported with local anesthetic use.

Full Prescribing Information is available at www.EXPAREL.com.

For more information, please visit www.EXPAREL.com or call 1-855-793-9727.